

## **BIBLIOGRAPHY**

### **General Adverse Drug Events**

Bates DW, Cullen DJ, et al., Incidence of adverse drug events and potential adverse drug events: implications for prevention. JAMA. 1995;274(1):29-34.

Bates DW, Spell N, et al, , The costs of adverse drug events in hospitalized patients. JAMA 1997;277(4):307-11

Forster AJ, Murff HJ, Peterson JF, Gandhi TK, Bates DW. The incidence and severity of adverse events affecting patients after discharge from the hospital. Ann Intern Med. 2003;138:161-167

Leape LL, Bates DW, Cullen DJ et al., Systems analysis of adverse drug events, JAMA. 1995;274:3543.

### **Medication History: Problems and Accuracy**

Andersen SE, Pedersen AB, Bach KF, Medication history on internal medicine wards: assessment of extra information collected from second drug interviews and GP lists. Pharmacoepidemiol Drug Saf. 2003;12(6):491-8

Bedell SE, Jabbour S, Goldberg R, Glaser H, Gobble S, Young-Xu, Y, Graboys TB and Ravid S. Discrepancies in the use of medications: their extent and predictors in an outpatient practice. Arch Intern Med. 2000;160:2129-2134

Beers MH, Munekata M and Storrie M. The accuracy of medication histories in the hospital medical records of elderly persons. J Am Geriatr Soc. 1990;38:1183-1187

Bobb A, Gleason K, Husch M, Feinglass J, Yarnold PR, and Noskin GA. The epidemiology of prescribing errors: the potential impact of computerized prescriber order entry. Arch Intern Med. 2004;164:785-792

Jacobson J. Ensuring continuity of care and accuracy of patients' medication history on hospital admission. Am J Health-Syst Pharm. 2002;59:1054-1055.

Lau HS, Florax C, Porsius AJ and de Boer, A. The completeness of medication histories in hospital medical records of patients admitted to general internal medicine wards. BrJ Clin Pharmacol. 2000;49:597-603.

Leister KA, Edwards WA, Christensen DB and Clarke H. A comparison of patient drug regimens as viewed by the physician, pharmacist and patient. *Med Care*. 1981; 19:658-64

.Lesar TS, Briceland L, Stein DS. Factors related to errors in medication prescribing." *JAMA*. 1997;277(4):312

Manley HJ, Drayer DK, McClaran M, Bender W, and Muther RS. Drug record discrepancies in an outpatient electronic medical record: frequency, type, and potential impact on patient care at a hemodialysis center. *Pharmacotherapy*. 2003;23:231-239.

Price D, Cooke J, Singleton S and Feely M. Doctors' unawareness of the drugs their patients are taking: a major cause of overprescribing? *BrMed J*. 1986;292:99-100.

Rozich JD and Resar RK. Medication safety: one organization's approach to the challenge. *J Clin Outcomes Manage*. 2001;8:27-34.

### **Medication Reconciliation**

Barnsteiner JH. Medication Reconciliation: Transfer of medication information across settings-keeping it free from error. *Am J Nurs*. 2005;105(3 Suppl):31-6

Gleason KM, Groszek JM, Sullivan C, Rooney D, Barnard C, and Noskin GA. Reconciliation of discrepancies in medication histories and admission orders of newly hospitalized patients. *Am J Health-Syst Pharm*. 2004;61:1689-1695.

Nickerson A, MacKinnon NJ, Roberts N, Saulnier L. Drug-therapy problems, inconsistencies and omissions identified during a medication reconciliation and seamless care service. *Healthc Q*. 2005;8

Pronovost P, Weast B, Schwarz M, Wyskiel RM, Prow D, Milanovich SN, Berenholtz S, Dorman T and Lopsett P. Medication reconciliation: a practical tool to reduce the risk of medication errors. *Journal of Critical Care*. 2003;18:201-205.

Provonost P, Weast W, et al. Medication reconciliation: a practical tool to reduce the risk of medication errors." *J Crit Care* 2003;18(4):201-5.

Rodehaver C, Fearing D. Medication reconciliation in acute care: ensuring an accurate drug regimen on admission and discharge. *Jt Comm J Qual Patient Saf*. 2005;31:406-13.

Rozich JD, Howard RM, Justeson JM, Macken PD, Lindsay ME and Resar RK. Standardization as a mechanism to improve safety in healthcare. *Jt Comm J Qua/ Saf*. 2004;30:5-14.

### **Pharmacists (Medication History – Adverse Events)**

Badowski SA, Rosenbloom D, and Dawson PH. Clinical importance of pharmacist-obtained medication histories using a validated questionnaire. *Am J Hosp Pharm.* 1984; 41:731-732

Bond CA, Raehl CL, and Franke T. Clinical pharmacy services, hospital pharmacy staffing and medication errors in United States hospitals. *Pharmacotherapy.* 2002; 22:134-147

Chapman NRM, Fotis MA, Yarnold PR and Gheorghiade M. Pharmacist interventions to improve the management of coronary artery disease. *Am J Health-Syst Pharm.* 2004;61:2672-2678

Jameson JP and Van Noord GR. Pharmacotherapy consultation on polypharmacy patients in ambulatory care. *Ann Pharmacother.* 2001;35:835-940

Kucukarslan SN, Peters M, Mlynarek M, and Nafziger DA. Pharmacists on rounding teams reduce preventable adverse drug events in hospital general medicine units. *Arch Intern Med.* 2003;163:2014-2018

Leape LL, Cullen DJ, Clapp MD, Burdick E, Demonaco HJ, Erickson JI and Bates DW. Pharmacist participation on physician rounds and adverse drug events in the intensive care unit. *JAMA.* 1999;282:267-270

Nester TM and Hale LS. Effectiveness of a pharmacist-acquired medication history in promoting patient safety. *Am J Health-Syst Pharm.* 2002;59:2221-2225.

Scarsi KK, Fotis MA and Noskin GA. Pharmacist participation in medical rounds reduces medication errors. *Am J Health-Syst Pharm.* 2002;59:2089-2092